

**bs-3136R****[ Primary Antibody ]****phospho-FLG (Tyr766) Rabbit pAb****Bioss**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —**


<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> <b>WB</b> (1:500-2000) <b>IHC-P</b> (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500)  <b>Reactivity:</b> Human, Mouse, Rat (predicted: Pig, Cow, Chicken, Dog, Horse)  <b>Predicted MW.:</b> 88 kDa  <b>Subcellular Location:</b> Cell membrane ,Cytoplasm ,Nucleus
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 2260	<b>SWISS:</b> P11362	
<b>Target:</b> FLG (Tyr766)		
<b>Immunogen:</b> KLH conjugated Synthesised phosphopeptide derived from human FGFR1 around the phosphorylation site of Tyr766: QE(p-Y)LD. < Cytoplasmic >		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> Fibroblast growth factors (FGFs) produce mitogenic and angiogenic effects in target cells by signaling through the cellular surface tyrosine kinase receptors. There are four members of the FGF receptor family: FGFR-1 (flg), FGFR-2 (bek, KGFR), FGFR-3 and FGFR-4. Each receptor contains an extracellular ligand binding domain, a transmembrane region and a cytoplasmic kinase domain (1). Following ligand binding and dimerization, the receptors are phosphorylated at specific tyrosine residues (2). Seven tyrosine residues in the cytoplasmic tail of FGFR-1 can be phosphorylated: Tyr463, Tyr583, Tyr585, Tyr653, Tyr654, Tyr730 and Tyr766. Tyrosine 653 and 654 are important for catalytic activity of the activated FGFR and are essential for signaling (3). The other phosphorylated tyrosine residues may provide docking sites for downstream signaling components such as Crk and PLCgamma.		

**— VALIDATION IMAGES —**


Sample: Cerebrum (Mouse) Lysate at 40 ug  
Primary: Anti-Phospho-FLG (Tyr766) (bs-3136R)  
at 1/300 dilution Secondary: IRDye800CW Goat  
Anti-Rabbit IgG at 1/20000 dilution Predicted  
band size: 88 kD Observed band size: 72 kD



Paraformaldehyde-fixed, paraffin embedded  
(human laryngeal carcinoma); Antigen retrieval  
by boiling in sodium citrate buffer (pH6.0) for  
15min; Block endogenous peroxidase by 3%  
hydrogen peroxide for 20 minutes; Blocking  
buffer (normal goat serum) at 37°C for 30min;  
Antibody incubation with (Phospho-FLG (Tyr766)  
) Polyclonal Antibody, Unconjugated (bs-3136R)  
at 1:200 overnight at 4°C, followed by operating  
according to SP Kit(Rabbit) (sp-0023)  
instructionsand DAB staining.



Paraformaldehyde-fixed, paraffin embedded  
(mouse brain); Antigen retrieval by boiling in  
sodium citrate buffer (pH6.0) for 15min; Block  
endogenous peroxidase by 3% hydrogen  
peroxide for 20 minutes; Blocking buffer (normal  
goat serum) at 37°C for 30min; Antibody  
incubation with (Phospho-FLG (Tyr766) )  
Polyclonal Antibody, Unconjugated (bs-3136R)  
at 1:200 overnight at 4°C, followed by operating  
according to SP Kit(Rabbit) (sp-0023)  
instructionsand DAB staining.



Tissue/cell: rat brain tissue;4%  
Paraformaldehyde-fixed and paraffin-  
embedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min;  
Blocking buffer (normal goat serum,C-0005) at 37℃ for 20 min; Incubation: Anti- p-FGFR1 Polyclonal Antibody, Unconjugated(bs-3136R) 1:200, overnight at 4℃; The secondary antibody was Goat Anti-Rabbit IgG, Cy3 conjugated(bs-0295G-Cy3)used at 1:200 dilution for 40 minutes at 37℃.  
DAPI(5ug/ml,blue,C-0033) was used to stain the cell nuclei