## bs-1642R

## [ Primary Antibody ]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

# phospho-FAK (Ser732) Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**GeneID: 5747 SWISS:** Q05397

Target: FAK (Ser732)

**Immunogen:** KLH conjugated Synthesised phosphopeptide derived from human

FAK around the phosphorylation site of Tyr732: YP(p-S)PQ.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** Non-receptor protein-tyrosine kinase implicated in signaling pathways involved in cell motility, proliferation and apoptosis. Activated by tyrosine-phosphorylation in response to either integrin clustering induced by cell adhesion or antibody crosslinking, or via G-protein coupled receptor (GPCR) occupancy by ligands such as bombesin or lysophosphatidic acid, or via LDL receptor occupancy. Plays a potential role in oncogenic transformations resulting in increased kinase activity. [SUBCELLULAR LOCATION] Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Note=Constituent of focal adhesions.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:100-500) Flow-Cyt (1µg /test) ICC/IF (1:100-500)

Reactivity: Human, Mouse


(predicted: Rat)

Predicted MW.: 116 kDa


Subcellular Cell membrane, Cytoplasm

Location: , Nucleus


## **VALIDATION IMAGES**




Paraformaldehyde-fixed, paraffin embedded (mouse brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (p-FAK(Ser732)) Polyclonal Antibody, Unconjugated (bs-1642R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



A549 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (phospho-FAK (Ser732)) polyclonal Antibody, Unconjugated (bs-1642R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.



Blank control (blue line): Hep G2 (blue). Primary Antibody (green line): Rabbit Anti-phospho-FAK(Ser732) antibody (bs-1642R) Dilution: 1µg /10^6 cells: Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody (white blue line): Goat anti-rabbit IgG-PE Dilution: 1µg /test. Protocol The cells were fixed with 70% ethanol (Overmight at 4°C) and then permeabilized with 90% ice-cold methanol for 30 min at -20°C.Cells stained with Primary Antibody for 30 min at room temperature. The cells were then incubated in 1 X PBS/2%BSA/10% goat serum to block nonspecific protein-protein interactions followed by the antibody for 15 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



Blank control:A549. Primary Antibody (green line): Rabbit Anti-phospho-FAK (Ser732) antibody (bs-1642R) Dilution: 1ug/Test; Secondary Antibody (white blue line): Goat antirabbit IgG-AF488 Dilution: 0.5ug/Test. Isotype control (orange line): Normal Rabbit IgG Protocol The cells were incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

### - SELECTED CITATIONS -

- [IF=16] Jing Zhang. et al. Robust and Regular Micronano Binary Texture on the Complex Curved Surface for Enhanced Reendothelialization and Antithrombotic Performance. ACS NANO. 2025;XXXX(XXX):XXX-XXX IF,FC;Human. 39813732
- [IF=5.673] Bott CJ et al. Nestin Selectively Facilitates the Phosphorylation of the Lissencephaly-Linked Protein Doublecortin (DCX) by cdk5/p35 to Regulate Growth Cone Morphology and Sema3a Sensitivity in Developing Neurons. J Neurosci. 2020 May 6;40(19):3720-3740. WB ;mouse. 32273484