

bs-34096R**[Primary Antibody]****NMDAR2B Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:200-1000) ICC/IF (1:50-200)
Clonality: Polyclonal		
GeneID: 2904	SWISS: Q13224	Reactivity: Human (predicted: Mouse, Rat)
Target: NMDAR2B		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Predicted MW.: 166 kDa
Background: N-methyl-D-aspartate (NMDA) receptors are a class of ionotropic glutamate receptors. NMDA receptor channel has been shown to be involved in long-term potentiation, an activity-dependent increase in the efficiency of synaptic transmission thought to underlie certain kinds of memory and learning. NMDA receptor channels are heteromers composed of three different subunits: NR1 (GRIN1), NR2 (GRIN2A, GRIN2B, GRIN2C, or GRIN2D) and NR3 (GRIN3A or GRIN3B). The NR2 subunit acts as the agonist binding site for glutamate. This receptor is the predominant excitatory neurotransmitter receptor in the mammalian brain. [provided by RefSeq, Jul 2008].		Subcellular Location: Cell membrane

— VALIDATION IMAGES —

4% Paraformaldehyde-fixed SH-SY5Y (H) cell;
Triton X-100 at r.t. for 20 min; Antibody
incubation with (NMDAR2B) polyclonal
Antibody, unconjugated (bs-34096R) 1:100, 90
min at 37°C; followed by conjugated Goat Anti-
Rabbit IgG antibody (green, bs-40295G-FITC) at
37°C for 90 min, DAPI (blue, C02-04002) was used
to stain the cell nuclei. PBS instead of the
primary antibody was used as the blank control.