

bs-4662R**[Primary Antibody]****phospho-TCF3 (Ser39) Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		IHC-P (1:100-500)
GeneID: 6929	SWISS: P15923	IHC-F (1:100-500)
Target: phospho-TCF3 (Ser39)		IF (1:100-500)
Immunogen: KLH conjugated synthesised phosphopeptide derived from human TCF3 around the phosphorylation site of Ser39: PA(p-S)LA.		
Purification: affinity purified by Protein A		Reactivity: Mouse, Rat (predicted: Human, Chicken, Dog)
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Predicted MW.: 67 kDa
Background: This gene encodes a member of the E protein (class I) family of helix-loop-helix transcription factors. E proteins activate transcription by binding to regulatory E-box sequences on target genes as heterodimers or homodimers, and are inhibited by heterodimerization with inhibitor of DNA-binding (class IV) helix-loop-helix proteins. E proteins play a critical role in lymphopoiesis, and the encoded protein is required for B and T lymphocyte development. Deletion of this gene or diminished activity of the encoded protein may play a role in lymphoid malignancies. This gene is also involved in several chromosomal translocations that are associated with lymphoid malignancies including pre-B-cell acute lymphoblastic leukemia (t(1;19), with PBX1), childhood leukemia (t(19;19), with TFPT) and acute leukemia (t(12;19), with ZNF384). Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene, and a pseudogene of this gene is located on the short arm of chromosome 9. [provided by RefSeq, Sep 2011]		Subcellular Location: Nucleus

— VALIDATION IMAGES —

Sample: Embryo (Mouse) Lysate at 40 ug Brain
(Rat) Lysate at 40 ug Primary: Anti-phospho-TCF3 (Ser39) (bs-4662R) at 1/300 dilution
Secondary: HRP conjugated Goat-Anti-rabbit IgG (bs-0295G-HRP) at 1/5000 dilution Predicted band size: 67 kD Observed band size: 68 kD



Tissue/cell: mouse embryo tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-phospho-TCF3/E2A/E47(Ser39) Polyclonal Antibody, Unconjugated(bs-4662R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

— SELECTED CITATIONS —

- **[IF=3.382]** Das D et al. Kaempferol inhibits extra-synaptic NMDAR mediated downregulation of TRK β in rat hippocampus during hypoxia. (2018) Neuroscience.Sep 26;392:77-91. WB ;Rat. 30266684