bsm-52289R

— DATASHEET –

[Primary Antibody]

Caspase-3 Recombinant Rabbit mAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Host: Rabbit Isotype: IgG Applications: WB (1:2000-10000) **Clonality:** Recombinant CloneNo.: 5A3 GenelD: 836 SWISS: P42574 Target: Caspase-3 Immunogen: A synthesized peptide derived from human Caspase 3: 52-120. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: The caspase family of cysteine proteases play a key role in apoptosis. Caspase 3 is the most extensively studied apoptotic protein among caspase family members. Caspase 3 is synthesized as inactive pro enzyme that is processed in cells undergoing apoptosis by self proteolysis and/or cleavage by other upstream proteases (e.g. Caspases 8, 9 and 10). The processed form of Caspase 3 consists of large (17kDa) and small (12kDa) subunits which associate to form an active enzyme. Caspase 3 is cleaved at Asp28 Ser29 and Asp175 Ser176. The active Caspase 3 proteolytically cleaves and activates other caspases (e.g. Caspases 6, 7 and 9), as well as relevant targets in the cells (e.g. PARP and DFF). Alternative splicing of this gene results in two transcript variants which encode the same protein. In immunohistochemical studies Caspase 3 expression has been shown to be widespread but not present in all cell types (e.g. commonly reported in epithelial cells of skin, renal proximal tubules and collecting ducts). Differences in the level of Caspase 3 have been reported in cells of short lived nature (eg germinal centre B cells) and those that are long lived (eg mantle zone B cells). Caspase 3 is the predominant caspase involved in the cleavage of amyloid beta 4A precursor

protein, which is associated with neuronal death in Alzheimer's

IHC-P (1:100-500) **IHC-F** (1:100-500) IF (1:100-500)

Reactivity: Human

Predicted 32 kDa MW.:

Subcellular Location: Cytoplasm

— VALIDATION IMAGES



disease.

25 ug total protein per lane of various lysates (see on figure) probed with Caspase-3 monoclonal antibody, unconjugated (bsm-52289R) at 1:3000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Human Spleen; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Caspase-3 Polyclonal Antibody, Unconjugated (bsm-52289R) at 1:200 overnight at 4°C, followed by conjugation to the SP Kit (Rabbit, SP-0023) and DAB (C-0010) staining



Paraformaldehyde-fixed, paraffin embedded Human Lymph; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Caspase-3 Polyclonal Antibody, Unconjugated (bsm-52289R) at 1:200 overnight at 4°C, followed by conjugation to the SP Kit (Rabbit, SP-0023) and DAB (C-0010) staining.

– SELECTED CITATIONS —

• [IF=2.3] Shiman Guo. et al. The effects of low ambient temperature on steroidogenesis and mitochondrial functions in

the testes of wild ground squirrels (Spermophilus dauricus). COMP BIOCHEM PHYS A. 2024 Jan;:111585 IHC ;Squirrels. 38228267