

bsm-52566R**[Primary Antibody]**

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

Caspase-9 Recombinant Rabbit mAb


DATASHEET

Host: Rabbit	Isotype: IgG	Applications: WB (1:1000-2000) IHC-P (1:50-200) IHC-F (1:50-200) IF (1:50-200)
Clonality: Recombinant	CloneNo.: 1D7	
GeneID: 842	SWISS: P55211	
Target: Caspase-9		
Immunogen: A synthesized peptide derived from human Caspase 9: 100-300.		
Purification: affinity purified by Protein A		Reactivity: Human, Mouse
Concentration: 1mg/ml		Predicted MW.: 35/50 kDa Subcellular Location: Cytoplasm
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Caspase 9 (also known as ICE like apoptotic protease 6 (ICE LAP6), apoptotic protease Mch6, and apoptotic protease activating factor 3 (Apaf3)) is a member of the peptidase family C14 that contains a CARD domain. This caspase is active as a heterotetramer and has been reported to have two isoforms. ProCaspase 9 has been reported to be approximately 47 kD. This caspase is present in the cytosol and, upon activation, translocates to the mitochondria. Caspase 9 is involved in the caspase activation cascade responsible for apoptosis execution and cleaves/activates Caspase 3 and Caspase 6. Caspase 9 is inhibited by the dominant negative isoform, BclXL, cIAP1, cIAP2, XIAP, and Livin. This caspase becomes activated when recruited to Apaf1/cytochrome c complex, and following cleavage by Apaf1, granzyme B, Caspase 3, possibly Caspase 8 and Caspase 10 into large p37 and small p10 subunits. Caspase 9 interacts with BIRC7 and has been shown to cleave PARP and vimentin.		


VALIDATION IMAGES



25 ug total protein per lane of various lysates (see on figure) probed with Caspase-9 monoclonal antibody, unconjugated (bsm-52566R) at 1:2000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Human Colon Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Caspase-9 Monoclonal Antibody, Unconjugated (bsm-52566R) at 1:200 overnight at 4°C, followed by conjugation to the SP Kit (Rabbit, SP-0023) and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Cervical Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Caspase-9 Monoclonal Antibody, Unconjugated (bsm-52566R) at 1:200 overnight at 4°C, followed by conjugation to the SP Kit (Rabbit, SP-0023) and DAB (C-0010) staining.

SELECTED CITATIONS

- **[IF=4.5]** Gan Rao. et al. Arsenic and polystyrene-nano plastics co-exposure induced testicular toxicity: Triggers oxidative stress and promotes apoptosis and inflammation in mice. ENVIRON TOXICOL. 2023 Sep; WB ;Mouse. 37705229
- **[IF=3.9]** Yuanliang Li. et al. Mn2O3 NPs-induced liver injury is potentially associated with gut microbiota dysbiosis in broiler chicken. FOOD CHEM TOXICOL. 2025 Aug;202:115487 WB ;Feather chicken. 40288515
- **[IF=2.817]** Qing Jin. et al. 125 I seeds irradiation inhibits tumor growth and induces apoptosis by Ki-67, P21, survivin,

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

livin and caspase-9 expression in lung carcinoma xenografts. Radiat Oncol. 2020 Dec;15(1):1-10 IHC ;Human. 33059701