## bs-19764R

## [ Primary Antibody ]

## **RBM16** Rabbit pAb




www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET		400-901-9800
Host: Rabbit	<b>lsotype:</b> lgG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal		IHC-F (1:100-500) IF (1:100-500)
GenelD: 22828	SWISS: Q9UPN6	
Target: RBM16		<b>Reactivity:</b> Rat (predicted: Human, Mouse, Rabbit)
Immunogen: KLH conjugated syn 1151-1271/1271.	thetic peptide derived from human RBM16	5:
Purification: affinity purified by Protein A		Predicted MW.: <sup>140 kDa</sup>
Concentration: 1mg/ml		MW.: 2 10 100
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: Nucleus

## — VALIDATION IMAGES -

Paraformaldehyde-fixed, paraffin embedded (Rat kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RBM16) Polyclonal Antibody, Unconjugated (bs-19764R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat colon); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RBM16) Polyclonal Antibody, Unconjugated (bs-19764R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.