bs-2197R

[Primary Antibody]

Duck hepatitis A virus polyprotein Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

| – DATASHEET – | | 400-901-9800 |
|--|---|---|
| Host: Rabbit Clonality: Polyclonal | lsotype: IgG | Applications: WB (1:500-2000) IHC-P (1:100-500) |
| Target: Duck hepatitis A virus polyprotein Immunogen: MAP peptide derived from polyprotein Duck hepatitis A virus (strain | | IHC-F (1:100-500) IF (1:100-500) ELISA (1:5000-10000) Reactivity: (predicted: Duck hepatitis A virus) |
| Duck/Taiwan/03D/2003): 851-950/2249. Purification: affinity purified by Protein A | | |
| Concentration: 1mg/ml | | |
| Glycerol. | n 1% BSA, 0.02% Proclin300 and 50% at -20°C for one year. Avoid repeated | Predicted MW.: 252 kDa |
| Background: Duck hepatitis is caused by the enteroviruses DHV-1 and DHV-3. It is a fatal disease of ducklings causing opisthotonus and hepatitis. Duck hepatitis virus, DHV, type 1 (DHV-1), type 1a (DHV-1a), type 2 (DHV-2), and type 3 (DHV-3) are the four types of viruses that can cause disease in ducks. The pathology and pathogenicity between duck hepatitis virus type 1 and type 1a is, in essence, the same as one another. | | |