

bs-2653R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

SPHK2 Rabbit pAb**— DATASHEET —**


Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ELISA (1:5000-10000) Reactivity: Human, Mouse Predicted MW.: 72 kDa Subcellular Location: Cell membrane ,Cytoplasm ,Nucleus
Clonality: Polyclonal		
GeneID: 56848	SWISS: Q9NRA0	
Target: SPHK2		
Immunogen: KLH conjugated synthetic peptide derived from human SPHK2: 101-200/654.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: This gene encodes a kinase that phosphorylates sphingosine into sphingosine-1-phosphate, which is involved in cell differentiation, motility, and apoptosis. The encoded protein plays a role in maintaining cellular levels of sphingosine-1-phosphate. The gene product also enhances apoptosis in different cell types and suppresses cellular proliferation. In mast cells, the encoded protein is necessary for influx of calcium, protein kinase C activation, and cytokine production and degranulation. Alternative splicing results in multiple transcript variants. [provided by RefSeq].		

— VALIDATION IMAGES —

25 ug total protein per lane of various lysates (see on figure) probed with SPHK2 polyclonal antibody, unconjugated (bs-2653R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Human Small Intestine; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with SPHK2 Polyclonal Antibody, Unconjugated (bs-2653R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with SPHK2 Polyclonal Antibody, Unconjugated (bs-2653R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.

— SELECTED CITATIONS —

- **[IF=15.2]** Lovren, Fina, et al. "MicroRNA-145 Targeted Therapy Reduces Atherosclerosis." *Circulation* 126.11 suppl 1 (2012): S81-S90. WB ;="Mouse". 22965997
- **[IF=5.853]** Mili Minaduola. et al. The circadian clock sets a spatial-temporal window for recent thymic emigrants. *IMMUNOL CELL BIOL.* 2022 Sep;; IF ;MOUSE. 36030488
- **[IF=4.29]** Tran, Hai B., et al. "Cigarette smoke inhibits efferocytosis via deregulation of sphingosine kinase signaling: reversal with exogenous S1P and the S1P analogue FTY720." *Journal of Leukocyte Biology* (2016): jlb-3A1015. Other ;="Human". 26792820
- **[IF=3.53]** Tani, Kohsuke, Keiji Tabuchi, and Akira Hara. "Hair Cell Loss Induced by Sphingosine and a Sphingosine

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Kinase Inhibitor in the Rat Cochlea."Neurotoxicity research 29.1 (2016): 35-46. WB,ICC ;="Rat" . 26472207

- **[IF=2.71]** Zhang, Ruxin, et al. "Hypoxic preconditioning protects cardiomyocytes against hypoxia/reoxygenation-induced cell apoptosis via sphingosine kinase 2 and FAK/AKT pathway." Experimental and Molecular Pathology (2015). WB ;="Rat" . 26621495