[Primary Antibody]

phospho-HSL (Ser660) Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET –––––		400-901-9800
Host: Rabbit	lsotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		IHC-P (1:100-500) IHC-F (1:100-500)
GenelD: 16890	SWISS: P54310	IF (1:100-500)
Target: HSL (Ser660)		ELISA (1:5000-10000)
Immunogen: KLH conjugated synthesised phosphopeptide derived from mouse HSL around the phosphorylation site of Ser660: SK(p-S)HE.		
Purification: affinity purified by P	rotein A	
Concentration: 1mg/ml		Predicted
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.		6 Predicted MW.: ^{83 kDa}
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: ^{Cell} membrane ,Cytoplasm
Background: HSL/LIPE is found in adipose tissue and heart, where it primarily hydrolyzes stored triglycerides to free fatty acids. It is also found in steroidogenic tissues, where it principally converts cholesteryl esters to free cholesterol for steroid hormone production. There are two named isoforms.		found in eryl

- SELECTED CITATIONS ------

- [IF=4.8] Huiling Hu. et al. Sestrin2 in POMC neurons modulates energy balance and obesity related metabolic disorders via mTOR signaling. J NUTR BIOCHEM. 2024 Nov;133:109703 WB ;Mouse. 39025457
- [IF=4.486] Chengqian Li. et al. Mutation of the novel acetylation site at K414R of BECN1 is involved in adipocyte differentiation and lipolysis. 2021 Jun 04 WB ;Human. 34085745