## bs-15186R

## [ Primary Antibody ]

## BIOSS ANTIBODIES www.bioss.com.cn

## C4a+C4b Rabbit pAb

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

**Clonality:** Polyclonal

**GenelD:** 100293534 **SWISS:** P0C0L5

Target: C4a+C4b

Immunogen: KLH conjugated synthetic peptide derived from human C4a and

C4b Complement C4 beta chain: 21-120/1744.

**Purification:** affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** This gene encodes the acidic form of complement factor 4, part of

the classical activation pathway. The protein is expressed as a single chain precursor which is proteolytically cleaved into a trimer of alpha, beta, and gamma chains prior to secretion. The trimer provides a surface for interaction between the antigen-antibody complex and other complement components. The alpha chain may be cleaved to release C4 anaphylatoxin, a mediator of local inflammation. Deficiency of this protein is associated with systemic lupus erythematosus and type I diabetes mellitus. This gene localizes to the major histocompatibility complex (MHC) class III region on chromosome 6. Varying haplotypes of this gene cluster exist, such that individuals may have 1, 2, or 3 copies of this gene. Two transcript variants encoding different isoforms have been

found for this gene. [provided by RefSeq, Nov 2011].

**Applications: ELISA** (1:5000-10000)

Reactivity: (predicted: Human)

Predicted 72/190 kDa

Subcellular Extracellular matrix, Cell

Location: membrane