

bsm-41222M**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

Lactoferrin Mouse mAb**— DATASHEET —**

Host: Mouse	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:400-800) IF (1:100-500) Reactivity: Human Predicted MW.: 77 kDa Subcellular Location: Secreted ,Cytoplasm ,Nucleus
Clonality: Monoclonal	CloneNo.: 4C1	
GeneID: 4057	SWISS: P02788	
Target: Lactoferrin		
Purification: affinity purified by Protein G		
Concentration: 1mg/ml		
Storage: Size : 50ul/100ul/200ul 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Size : 200ug (PBS only) 0.01M PBS Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Transferrins are iron binding transport proteins which can bind two Fe(3+) ions in association with the binding of an anion, usually bicarbonate. Lactotransferrin has antimicrobial activity which depends on the extracellular cation concentration. Lactoferroxins A, B and C have opioid antagonist activity. Lactoferroxin A shows preference for mu-receptors, while lactoferroxin B and C have somewhat higher degrees of preference for kappa-receptors than for mu-receptors. The lactotransferrin transferrin-like domain 1 functions as a serine protease of the peptidase S60 family that cuts arginine rich regions. This function contributes to the antimicrobial activity.		

— VALIDATION IMAGES —

Sample: Lane 1: Human Lactoferrin(bs-41222P)
 Primary: Anti-Lactoferrin (bsm-41222M) at 1/1000 dilution
 Secondary: IRDye800CW Goat Anti-Mouse IgG at 1/20000 dilution
 Predicted band size: 77 kDa
 Observed band size: 77 kDa



Paraformaldehyde-fixed, paraffin embedded (human spleen); Antigen retrieval by boiling in E sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (Lactoferrin) Monoclonal Antibody, Unconjugated (bsm-41222M) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructions and DAB staining.