
phospho-Tau (Ser416) Rabbit pAb

Catalog Number: bs-5473R

Target Protein: phospho-Tau (Ser416)

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), Flow-Cyt (1µg/Test)

Reactivity: Human, Mouse, Rat (predicted:Cow, Chicken, Dog, Horse)

Predicted MW: 52/79 kDa

Entrez Gene: 4137

Swiss Prot: P10636

Source: KLH conjugated Synthesised phosphopeptide derived from human Tau around the phosphorylation site of Ser416: TG(p-S)ID.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Tau proteins are important Promotes microtubule assembly and stability, and might be involved in the establishment and maintenance of neuronal polarity. The C-terminus binds axonal microtubules while the N-terminus binds neural plasma membrane components, suggesting that tau functions as a linker protein between both. Axonal polarity is predetermined by tau localization (in the neuronal cell) in the domain of the cell body defined by the centrosome. The short isoforms allow plasticity of the cytoskeleton whereas the longer isoforms may preferentially play a role in its stabilization. Tau proteins subcellular located in the axons of neurons, in the cytosol and in association with plasma membrane components. It expressed in neurons. PNS-tau is expressed in the peripheral nervous system while the others are expressed in the central nervous system.


VALIDATION IMAGES




Sample: Cerebrum (Mouse) Lysate at 40 ug Cerebrum (Rat) Lysate at 40 ug Primary: Anti-phospho-Tau (Ser416) (bs-5473R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 52/79 kD Observed band size: 52 kD



Sample: Lane 1: Cerebrum (Rat) Lysate at 40 ug Lane 2: Cerebrum (Mouse) Lysate at 40 ug Lane 3: Cerebellum (Rat) Lysate at 40 ug Lane 4: Cerebellum (Mouse) Lysate at 40 ug Primary: Anti-phospho-Tau (Ser416) (bs-5473R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 50-70 kD Observed band size: 55' 62kD



Sample: Lane 1: Cerebrum (Rat) Lysate at 40 ug Lane 2: Cerebrum (Mouse) Lysate at 40 ug Lane 3: Cerebellum (Rat) Lysate at 40 ug Lane 4: Cerebellum (Mouse) Lysate at 40 ug Primary: Anti-phospho-Tau (Ser416) (bs-5473R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 50-70 kD Observed band size: 55' 62kD



Blank control (blue line): MCF 7 (fixed with 70% methanol (Overnight at 4°C) and then permeabilized with 90% ice-cold methanol for 30 min on ice). Primary Antibody (green line): Rabbit Anti-phospho-Tau (Ser416) antibody (bs-5473R), Dilution: 1µg /10⁵ cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody (white blue line): Goat anti-rabbit IgG-FITC, Dilution: 1µg /test.