

bs-5473R**[Primary Antibody]****phospho-Tau (Ser416) Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —


Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) Flow-Cyt (1µg/Test)
Clonality: Polyclonal		
GeneID: 4137	SWISS: P10636	Reactivity: Human, Mouse, Rat (predicted: Cow, Chicken, Dog, Horse)
Target: Tau (Ser416)		Predicted MW.: 52/79 kDa
Immunogen: KLH conjugated Synthesised phosphopeptide derived from human Tau around the phosphorylation site of Ser416: TG(p-S)ID.		Subcellular Location: Cell membrane ,Cytoplasm
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Tau proteins are important Promotes microtubule assembly and stability, and might be involved in the establishment and maintenance of neuronal polarity. The C-terminus binds axonal microtubules while the N-terminus binds neural plasma membrane components, suggesting that tau functions as a linker protein between both. Axonal polarity is predetermined by tau localization (in the neuronal cell) in the domain of the cell body defined by the centrosome. The short isoforms allow plasticity of the cytoskeleton whereas the longer isoforms may preferentially play a role in its stabilization. Tau proteins subcellular located in the axons of neurons, in the cytosol and in association with plasma membrane components. It expressed in neurons. PNS-tau is expressed in the peripheral nervous system while the others are expressed in the central nervous system.		

— VALIDATION IMAGES —

Sample: Lane 1: Cerebrum (Rat) Lysate at 40 ug
Lane 2: Cerebrum (Mouse) Lysate at 40 ug Lane
3: Cerebellum (Rat) Lysate at 40 ug Lane 4:
Cerebellum (Mouse) Lysate at 40 ug Primary:
Anti-phospho-Tau (Ser416) (bs-5473R) at 1/1000
dilution Secondary: IRDye800CW Goat Anti-
Rabbit IgG at 1/20000 dilution Predicted band
size: 50-70 kD Observed band size: 55' 62kD



Sample: Lane 1: Cerebrum (Rat) Lysate at 40 ug
Lane 2: Cerebrum (Mouse) Lysate at 40 ug Lane
3: Cerebellum (Rat) Lysate at 40 ug Lane 4:
Cerebellum (Mouse) Lysate at 40 ug Primary:
Anti-phospho-Tau (Ser416) (bs-5473R) at 1/1000
dilution Secondary: IRDye800CW Goat Anti-
Rabbit IgG at 1/20000 dilution Predicted band
size: 50-70 kD Observed band size: 55' 62kD



Blank control (blue line): MCF 7 (fixed with 70% methanol (Overnight at 4°C) and then permeabilized with 90% ice-cold methanol for 30 min on ice). Primary Antibody (green line): Rabbit Anti-phospho-Tau (Ser416) antibody (bs-5473R), Dilution: 1µg / 10⁵ cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody (white blue line): Goat anti-rabbit IgG-FITC, Dilution: 1µg / test.