
phospho-P38 MAPK (Thr180) Rabbit pAb

Catalog Number: bs-5476R

Target Protein: phospho-P38 MAPK (Thr180)

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500), Flow-Cyt (1µg/Test)

Reactivity: Human, Mouse, Rat (predicted:Rabbit, Pig, Chicken, Dog, Horse)

Predicted MW: 41 kDa

Subcellular: Cytoplasm, Nucleus

Locations:

Entrez Gene: 1432

Swiss Prot: Q16539

Source: KLH conjugated Synthesised phosphopeptide derived from human MAPK14 around the phosphorylation site of Thr180: EM(p-T)G.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase is activated by various environmental stresses and proinflammatory cytokines. The activation requires its phosphorylation by MAP kinase kinases(MKKs), or its autophosphorylation triggered by the interaction of MAP3K7IP1/TAB1 protein with this kinase. The substrates of this kinase include transcription regulator ATF2, MEF2C, and MAX, cell cycle regulator CDC25B, and tumor suppressor p53, which suggest the roles of this kinase in stress related transcription and cell cycle regulation, as well as in genotoxic stress response. Four alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.


VALIDATION IMAGES




Sample: Muscle (Mouse) Lysate at 40 ug Muscle (Rat) Lysate at 40 ug Primary: Anti- phospho-P38 MAPK (Thr180) (bs-5476R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 41 kD Observed band size: 41 kD




Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (p-P38 MAPK (Thr180)) Polyclonal Antibody, Unconjugated (bs-5476R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-P38 MAPK (Thr180)) Polyclonal Antibody, Unconjugated (bs-5476R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (p-P38 MAPK (Thr180)) Polyclonal Antibody, Unconjugated (bs-5476R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control(blue): HeLa (fixed with 2% paraformaldehyde (10 min), then permeabilized with 0.3% Triton X-100 for 5 min at room temperature). Primary Antibody: Rabbit Anti-phospho-P38 MAPK (Thr180) antibody (bs-5476R) Dilution: 1 µg in 100 µL 1X PBS containing 0.5% BSA; Isotype Control Antibody: Rabbit IgG (orange), used under the same conditions; Secondary Antibody: Goat anti-rabbit IgG-FITC, Dilution: 1:200 in 1 X PBS containing 0.5% BSA.

PRODUCT SPECIFIC PUBLICATIONS

[IF=8.039] Yifan Zhu. et al. Discovery of Selective P2Y6R Antagonists with High Affinity and In Vivo Efficacy for Inflammatory Disease Therapy. J MED CHEM. 2023;XXX(XXX):XXX-XXX **WB ; Mouse** . 37078976

[IF=7.59] Muzhe Li. et al. STS load PCL- MECM based hydrogel hybrid scaffold promote meniscal regeneration via modulating macrophage phenotype polarization. BIOMATER SCI-UK. 2023 Jan;; **WB ; Rabbit** . 10.1039/D2BM00526C

[IF=7.9] Keyi Nong. et al. Potential effects and mechanism of flavonoids extract of Callicarpa nudiflora Hook on DSS-induced colitis in mice. PHYTOMEDICINE. 2024 Jun;128:155523 **WB ; Mouse** . 38489893

[IF=8.2] Xinyun Qin. et al. Regulation of the intestinal flora using polysaccharides from Callicarpa nudiflora Hook to alleviate ulcerative colitis and the molecular mechanisms involved. INT J BIOL MACROMOL. 2024 Feb;258:128887 **WB ; Mouse** . 38118262

[IF=8.025] Lihui Ni. et al. The combination of insulin and linezolid ameliorates Staphylococcus aureus pneumonia in individuals with diabetes via the TLR2/MAPKs/NLRP3 pathway. INT J BIOL MACROMOL. 2023 Jul;242:124750 **WB ; Mouse** . 37160172