bs-33016R

[Primary Antibody]

BIOSS ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn

Strep-Tag II Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Target: Strep-Tag II

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.


Background: The Strep-tag system is a method which allows the purification

and detection of proteins by affinity chromatography. The Streptag is a synthetic peptide consisting of eight amino acids (Trp-Ser-His-Pro-Gln-Phe-Glu-Lys). This peptide sequence exhibits intrinsic affinity towards Strep-Tactin, a specifically engineered streptavidin and can be N- or C- terminally fused to recombinant proteins. By exploiting the highly specific interaction, Strep-tagged proteins can be isolated in one step from crude cell lysates. Because the Strep-tag elutes under gentle, physiological conditions it is especially suited for generation of functional proteins.

VALIDATION IMAGES



Sample: Lane 1: Strep-Tagged Fusion Protein Overexpression E.coli Lysate (Cat#: bs-41403P) Lane 2: Negative control Primary: Anti-Strep-Tag II (bs-33016R) at 1/10000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 51 kD Observed band size: 51 kD



Sample: Lane 1: Strep-Tagged Fusion Protein Overexpression E.coli Lysate (Cat#: bs-41403P) at 2ug Lane 2: Strep-Tagged Fusion Protein Overexpression E.coli Lysate (Cat#: bs-41403P) at 4ug Primary: Anti-Strep-Tag II (bs-33016R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 51 kD Observed band size: 51 kD Reactivity: Species independent

Subcellular Secreted