
TCF3 Recombinant Rabbit mAb

Catalog Number: bsm-60695R

Target Protein: TCF3

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Recombinant

Clone No.: R4F10

Isotype: IgG

Applications: WB (1:500-1000), IHC-P (1:100-500), IHC-F (1:200-500), IF (1:50-100)

Reactivity: Human

Predicted MW: 67 kDa

Entrez Gene: 6929

Swiss Prot: P15923


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: This gene encodes a member of the E protein (class I) family of helix-loop-helix transcription factors. E proteins activate transcription by binding to regulatory E-box sequences on target genes as heterodimers or homodimers, and are inhibited by heterodimerization with inhibitor of DNA-binding (class IV) helix-loop-helix proteins. E proteins play a critical role in lymphopoiesis, and the encoded protein is required for B and T lymphocyte development. Deletion of this gene or diminished activity of the encoded protein may play a role in lymphoid malignancies. This gene is also involved in several chromosomal translocations that are associated with lymphoid malignancies including pre-B-cell acute lymphoblastic leukemia (t(1;19), with PBX1), childhood leukemia (t(19;19), with TFPT) and acute leukemia (t(12;19), with ZNF384). Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene, and a pseudogene of this gene is located on the short arm of chromosome 9. [provided by RefSeq, Sep 2011]

VALIDATION IMAGES



Blocking buffer: 5% NFDM/TBST Primary Ab dilution: 1:5000 Primary Ab incubation condition: 2 hours at room temperature Secondary Ab: Goat Anti-Rabbit IgG H&L (HRP) Lysate: 1: Daudi, 2: Ramos Protein loading quantity: 20 μ g Exposure time: 60 s Predicted MW: 68 kDa Observed MW: 68 kDa