

**bs-13471R****[ Primary Antibody ]****GNL2 Rabbit pAb****Bioss**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —**


<b>Host:</b> Rabbit <b>Clonality:</b> Polyclonal <b>GeneID:</b> 29889 <b>Target:</b> GNL2 <b>Immunogen:</b> KLH conjugated synthetic peptide derived from human GNL2: 1-100/731. <b>Purification:</b> affinity purified by Protein A <b>Concentration:</b> 1mg/ml <b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. <b>Background:</b> GNL2 is a nucleolar guanine-triphosphate binding protein that is ubiquitously expressed at low levels in almost all tissues. GNL2 is involved in the crucial process of trafficking proteins out of the nucleus. Specifically, it is a GTPase that interacts with the 60s preribosomal subunit in the nucleus and facilitates export of the subunit into the cytoplasm. GTPases are responsible for the hydrolysis of GTP by way of a protein region dubbed the G domain. GTPases are often involved in the translocating proteins through membranes gleaned energy for the activity by hydrolyzing GTP. GNL2 shares G domain homology and some functionality with nucleostemin (GNL3), another nuclear GTPase. Highest expression of GNL2 is found in testis.	<b>Isotype:</b> IgG  <b>SWISS:</b> Q13823	<b>Applications:</b> <b>WB</b> (1:500-2000) <b>IHC-P</b> (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500)
		<b>Reactivity:</b> Mouse, Rat (predicted: Human, Pig, Sheep, Cow, Dog, Horse)
		<b>Predicted MW.:</b> 84 kDa
		<b>Subcellular Location:</b> Nucleus

**— VALIDATION IMAGES —**

Sample: Muscle (Mouse) Lysate at 40 ug Primary:  
Anti- GNL2 (bs-13471R) at 1/300 dilution  
Secondary: IRDye800CW Goat Anti-Rabbit IgG at  
1/20000 dilution Predicted band size: 84 kD  
Observed band size: 84 kD



Tissue/cell: Rat testis tissue; 4%  
Paraformaldehyde-fixed and paraffin-  
embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block  
endogenous peroxidase by 3% Hydrogen  
peroxide for 30min; Blocking buffer (normal goat  
serum, C-0005) at 37°C for 20 min; Incubation:  
Anti-GNL2 Polyclonal Antibody,  
Unconjugated(bs-13471R) 1:500, overnight at  
4°C, followed by conjugation to the secondary  
antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: Mouse brain tissue; 4%  
Paraformaldehyde-fixed and paraffin-  
embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block  
endogenous peroxidase by 3% Hydrogen  
peroxide for 30min; Blocking buffer (normal goat  
serum, C-0005) at 37°C for 20 min; Incubation:  
Anti-GNL2 Polyclonal Antibody,  
Unconjugated(bs-13471R) 1:500, overnight at  
4°C, followed by conjugation to the secondary  
antibody(SP-0023) and DAB(C-0010) staining