

bs-19400R**[Primary Antibody]****GUCA2B Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000) Reactivity: (predicted: Human, Cow) Predicted MW.: 12 kDa Subcellular Location: Secreted
Clonality: Polyclonal		
GeneID: 2981	SWISS: Q16661	
Target: GUCA2B		
Immunogen: KLH conjugated synthetic peptide derived from human GUCA2B: 89-112/112.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: The family of guanylin regulatory peptides, including guanylin and uroguanylin, are strongly expressed in intestinal mucosa and regulate intestinal fluid secretion during digestion. Guanylins are also involved in acid neutralization and the regulation of membrane-bound guanylate cyclase signaling molecules. Guanylin and uroguanylin are secreted primarily in the stomach, intestine, and colon. Uroguanylin is an endogenous activator of intestinal guanylate cyclase. It is a paracrine and/or autocrine regulator of intestinal water and salt transport. Uroguanylin stimulates intestinal guanylate cyclase through the same receptor binding region as the heat-stable enterotoxins. Uroguanylin is involved in the regulation of intestinal fluid and electrolyte transport.		