


bsm-52978R**[Primary Antibody]****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

FGF21 Recombinant Rabbit mAb**— DATASHEET —****Host:** Rabbit**Isotype:** IgG**Clonality:** Recombinant**CloneNo.:** 7C4**GeneID:** 26291**SWISS:** Q9NSA1**Target:** FGF21**Immunogen:** A synthesized peptide derived from human FGF21: 5-61.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.**Background:** FGF21 is a growth factor that is a member of the FGF family. Proteins of this family play a central role during prenatal development and postnatal growth and regeneration of a variety of tissues, by promoting cellular proliferation and differentiation. FGF21, in the presence of betaKlotho as a protein cofactor, signals through the FGFR 1c and 4 receptors and stimulates insulin independent glucose uptake by adipocytes.**Applications:** **WB** (1:1000-5000)**IHC-P** (1:200-800)**IHC-F** (1:200-800)**IF** (1:200-800)**Reactivity:** Human, Mouse, Rat**Predicted**
MW.: 20 kDa**Subcellular**
Location: Secreted**— VALIDATION IMAGES —**

25 ug total protein per lane of various lysates (see on figure) probed with FGF21 monoclonal antibody, unconjugated (bsm-52978R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Human Spleen; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with FGF21 Monoclonal Antibody, Unconjugated(bsm-52978R) at 1:250 overnight at 4°C, followed by conjugation to the SP Kit(Rabbit, SP-0023) and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Rat Spleen; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with FGF21 Monoclonal Antibody, Unconjugated(bsm-52978R) at 1:250 overnight at 4°C, followed by conjugation to the SP Kit(Rabbit, SP-0023) and DAB (C-0010) staining.