


bsm-60605R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

CD19 Recombinant Rabbit mAb**— DATASHEET —****Host:** Rabbit**Isotype:** IgG**Clonality:** Recombinant**GeneID:** 930**SWISS:** P15391**Target:** CD19**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.**Background:** This gene encodes a member of the immunoglobulin gene superfamily. Expression of this cell surface protein is restricted to B cell lymphocytes. This protein is a reliable marker for pre-B cells but its expression diminishes during terminal B cell differentiation in antibody secreting plasma cells. The protein has two N-terminal extracellular Ig-like domains separated by a non-Ig-like domain, a hydrophobic transmembrane domain, and a large C-terminal cytoplasmic domain. This protein forms a complex with several membrane proteins including complement receptor type 2 (CD21) and tetraspanin (CD81) and this complex reduces the threshold for antigen-initiated B cell activation. Activation of this B-cell antigen receptor complex activates the phosphatidylinositol 3-kinase signalling pathway and the subsequent release of intracellular stores of calcium ions. This protein is a target of chimeric antigen receptor (CAR) T-cells used in the treatment of lymphoblastic leukemia. Mutations in this gene are associated with the disease common variable immunodeficiency 3 (CVID3) which results in a failure of B-cell differentiation and impaired secretion of immunoglobulins. CVID3 is characterized by hypogammaglobulinemia, an inability to mount an antibody response to antigen, and recurrent bacterial infections. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Jul 2020]**Applications:** **WB** (1:500-2000)
IHC-P (1:100-500)
IHC-F (1:400-800)
IF (1:100-500)
Flow-Cyt (1:20-50)
ICC/IF (1:20-50)**Reactivity:** Human**Predicted MW.:** 59 kDa**Subcellular Location:** Cell membrane**— VALIDATION IMAGES —**


Sample: Lane 1: Human Raji cell Lysates
 Primary: Anti-CD19 (bsm-60605R) at 1/1000
 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 59kDa Observed band size: 90kDa




Tissue: Human tonsil Section type: Formalin fixed & Paraffin -embedded section Retrieval method: High temperature and high pressure Retrieval buffer: Tris/EDTA buffer, pH 9.0 Primary ab dilution: 1:1000 Primary ab incubation condition: 1 hour at room temperature Secondary ab: SP Kit(Rabbit)(sp-0023) Counter stain: Hematoxylin (Blue) Comment: Color brown is the positive signal for bsm-60605R




Tissue: Human spleen Section type: Formalin fixed & Paraffin -embedded section Retrieval method: High temperature and high pressure Retrieval buffer: Tris/EDTA buffer, pH 9.0 Primary ab dilution: 1:1000 Primary ab incubation condition: 1 hour at room temperature Secondary ab: SP Kit(Rabbit) Counter stain: Hematoxylin (Blue) Comment: Color brown is the positive signal for bsm-60605R



Cell line: Raji Fixative: 4% Paraformaldehyde
 Permeabilization: 0.1% TritonX-100 Primary ab
 dilution: 1:50 Primary incubation condition: 4°C
 overnight Secondary ab: Goat Anti-Rabbit IgG
 Nuclear counter stain: DAPI (Blue) Comment:
 Color green is the positive signal for bsm-60605R



Cell line: Raji Fixative: 4% Paraformaldehyde
 Permeabilization: 90% Methanol Primary ab
 dilution: 1:100 Secondary ab: Goat anti Rabbit
 IgG Unlabelled control: The cell without
 incubation with primary antibody and secondary
 antibody (Black line). Isotype control: Rabbit
 monoclonal IgG (Blue line). Comment: Line red is
 the positive signal for bsm-60605R



Cell line: Ramos Fixative: 4% Paraformaldehyde
 Permeabilization: 90% Methanol Primary ab
 dilution: 1:100 Secondary ab: Goat anti Rabbit
 IgG Unlabelled control: The cell without
 incubation with primary antibody and secondary
 antibody (Black line). Isotype control: Rabbit
 monoclonal IgG (Blue line). Comment: Line red is
 the positive signal for bsm-60605R