

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

## **ABCA1** Recombinant Mouse mAb

Catalog Number: bsm-60239M

Target Protein: ABCA1
Concentration: 1mg/ml

Form: Liquid

Host: Mouse

Clonality: Recombinant

Isotype: IgG

Applications: WB (1:500-2000)

Reactivity: Mouse Predicted MW: 254 kDa

Entrez Gene: 19

Swiss Prot: 095477

Purification: affinity purified by Protein G

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: The membrane-associated protein encoded by this gene is a member of the superfamily of

ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intracellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the ABC1 subfamily. Members of the ABC1 subfamily comprise the only major ABC subfamily found exclusively in multicellular eukaryotes. With cholesterol as its substrate, this protein functions as a cholesteral efflux pump in the cellular lipid removal pathway. Mutations in both alleles of this gene cause Tangier disease and familial high-density lipoprotein (HDL)

deficiency. [provided by RefSeq, Sep 2019]

## **VALIDATION IMAGES**



Blocking buffer: 5% NFDM/TBST Primary ab dilution: 1:2000 Primary ab incubation condition: 2 hours at room temperature Lysate: Mouse liver Protein loading quantity:  $20~\mu g$  Exposure time: 180~S Predicted MW: 254~kDa Observed MW: 254~kDa

## PRODUCT SPECIFIC PUBLICATIONS

[IF=5.4] Chun-xia Nie. et al. Shenlian extract protected ox-LDL-loaded macrophages against ER stress by promoting LAL-LXR $\alpha$  mediated cholesterol flux. J ETHNOPHARMACOL. 2023 Jun;:116721 WB; MOUSE . 37315648

[IF=4.8] Haoqiang Chen. et al. IFIT2 mediates iron retention and cholesterol efflux in atherosclerosis. INT IMMUNOPHARMACOL. 2024 Dec;142:113131 WB,IF; Mouse . 39276454

[IF=3.3] Yueqi Cui. et al. The activation of liver X receptors in Madin-Darby bovine kidney cells and mice restricts infection by bovine viral diarrhea virus. VET MICROBIOL. 2023 Dec;:109948 WB; Bovine . 38113573