

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DAZL Rabbit pAb

Catalog Number: bs-12245R

Target Protein: DAZL
Concentration: 1mg/ml

Form: Liquid Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Mouse (predicted:Human, Rabbit, Pig, Sheep, Cow, Dog, Horse)

Predicted MW: 33 kDa

Subcellular Cytoplasm, Nucleus

Locations:

Entrez Gene: 1618 Swiss Prot: Q92904

Source: KLH conjugated synthetic peptide derived from human DAZL: 51-130/295.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: The DAZ (Deleted in AZoospermia) gene family encodes potential RNA binding proteins that

are expressed in prenatal and postnatal germ cells of males and females. The protein

encoded by this gene is localized to the nucleus and cytoplasm of fetal germ cells and to the

cytoplasm of developing oocytes. In the testis, this protein is localized to the nucleus of

spermatogonia but relocates to the cytoplasm during meiosis where it persists in


spermatids and spermatozoa. Transposition and amplification of this autosomal gene

during primate evolution gave rise to the DAZ gene cluster on the Y chromosome. Mutations in this gene have been linked to severe spermatogenic failure and infertility in males. Two

transcript variants encoding different isoforms have been found for this gene. [provided by

RefSeq, Jun 2010].

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (DAZL) Polyclonal Antibody, Unconjugated (bs-12245R) at 1:400 overnight at 4°C, followed by a conjugated Goat Anti-Rabbit IgG antibody (bs-0295G-Cy3) for 90 minutes, and DAPI for nuclei staining.

PRODUCT SPECIFIC PUBLICATIONS

[IF=4.2] Zhang, Weidong, et al. "Decrease in male mouse fertility by hydrogen sulfide and/or ammonia can Be inheritable." Chemosphere (2017). IHC; = "Mouse". 29202267

[IF=2.638] Li T et al. Gene expression patterns and protein cellular localization suggest a novel role for DAZL in developing Tibetan sheep testes. Gene. 2020 Mar 20;731:144335. WB,IHC,IF; Tibetan sheep ram . 31927007

[IF=3.4] Ünal Gülin Özdamar. et al. The beneficial effects of vortioxetine on BDNF, CREB, S100B, β amyloid, and glutamate NR2b receptors in chronic unpredictable mild stress model of depression. PSYCHOPHARMACOLOGY. 2023 Aug;:1-15 IHC; Rat. 37555927