

**bs-5842R****[ Primary Antibody ]****MIIP Rabbit pAb****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit <b>Clonality:</b> Polyclonal <b>GeneID:</b> 60672 <b>Target:</b> MIIP <b>Immunogen:</b> KLH conjugated synthetic peptide derived from human MIIP: 251-350/388. <b>Purification:</b> affinity purified by Protein A <b>Concentration:</b> 1mg/ml <b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. <b>Background:</b> MIIP has 3 SEG (segments of low compositional complexity) domains, an RGD motif, and several potential phosphorylation sites. The C-terminal region of IGFBP2 interacts with a central 44-amino acid sequence of MIIP. MIIP inhibits glioma cells invasion and down-regulates adhesion- and motility-associated genes such as NFKB2 and ICAM1. It exhibits opposing effects to IGFBP2 on cell invasion. There are 2 named isoforms due to alternative splicing. Isoform 1 is expressed in brain but underexpressed in glioma tissues, at protein level. Isoform 2 is not detected in normal organs, but is expressed in gliomas with increasing levels with glioma progression. On the contrary, at protein level, isoform 2 is not detected in gliomas, suggesting that this isoform is unstable in glioma cells. Isoform 2 is degraded by the ubiquitin-proteasome pathway.	<b>Isotype:</b> IgG <b>SWISS:</b> Q5JXC2 <b>Applications:</b> IHC-P (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500) <b>Reactivity:</b> Human (predicted: Mouse, Rat, Rabbit, Cow, Dog, Horse) <b>Predicted MW.:</b> 43 kDa <b>Subcellular Location:</b> Cytoplasm ,Nucleus
---	--

**— VALIDATION IMAGES —**

Paraformaldehyde-fixed, paraffin embedded (Human liver carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (MIIP) Polyclonal Antibody, Unconjugated (bs-5842R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.