

bsm-60582R**[Primary Antibody]**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

Inhibin alpha Recombinant Rabbit mAb


— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-1000) IHC-P (1:200-1:1000) IHC-F (1:50-100) IF (1:50-100) ICC/IF (1:50-100) Reactivity: Human Predicted MW.: 15 kDa Subcellular Location: Secreted
Clonality: Recombinant	CloneNo.: A8H6	
GeneID: 3623	SWISS: P05111	
Target: Inhibin alpha		
Immunogen: A synthesized peptide derived from human Inhibin alpha chain: 200-366/366.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: The inhibin alpha subunit joins either the beta A or beta B subunit to form a pituitary FSH secretion inhibitor. Inhibin has been shown to regulate gonadal stromal cell proliferation negatively and to have tumour-suppressor activity. In addition, serum levels of inhibin have been shown to reflect the size of granulosa-cell tumors and can therefore be used as a marker for primary as well as recurrent disease. However, in prostate cancer, expression of the inhibin alpha-subunit gene was suppressed and was not detectable in poorly differentiated tumor cells. Furthermore, because expression in gonadal and various extragonadal tissues may vary severalfold in a tissue-specific fashion, it is proposed that inhibin may be both a growth/differentiation factor and a hormone.		

— VALIDATION IMAGES —



Paraformaldehyde-fixed, paraffin embedded Human Adrenal Gland; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Inhibin alpha Monoclonal Antibody, Unconjugated(bsm-60582R) at 1:200 overnight at 4°C, followed by conjugation to the SP Kit(Rabbit, SP-0023) and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Testicles; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Inhibin alpha Monoclonal Antibody, Unconjugated(bsm-60582R) at 1:200 overnight at 4°C, followed by conjugation to the SP Kit(Rabbit, SP-0023) and DAB (C-0010) staining.