

bs-12262R**[Primary Antibody]****Phospho-TIE2 (Tyr1108) Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —


Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Human, Mouse, Rat (predicted: Rabbit, Pig, Sheep, Cow, Chicken, Dog, Horse) Predicted MW.: 124 kDa Subcellular Location: Secreted ,Extracellular matrix ,Cell membrane ,Cytoplasm
Clonality: Polyclonal		
GeneID: 7010	SWISS: Q02763	
Target: Phospho-TIE2 (Tyr1108)		
Immunogen: KLH conjugated Synthesised phosphopeptide derived from human TIE2 around the phosphorylation site of Tyr1108: TL(p-Y)EK.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: The TEK receptor tyrosine kinase is expressed almost exclusively in endothelial cells in mice, rats, and humans. This receptor possesses a unique extracellular domain containing 2 immunoglobulin-like loops separated by 3 epidermal growth factor-like repeats that are connected to 3 fibronectin type III-like repeats. The ligand for the receptor is angiopoietin-1. Defects in TEK are associated with inherited venous malformations; the TEK signaling pathway appears to be critical for endothelial cell-smooth muscle cell communication in venous morphogenesis. TEK is closely related to the TIE receptor tyrosine kinase.		

— VALIDATION IMAGES —

Sample: Lane 1: Lung (Rat) Lysate at 40 ug Lane 2: Kidney (Rat) Lysate at 40 ug Primary: Anti-Phospho-TIE2 (Tyr1108) (bs-12262R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 140 kD Observed band size: 140 kD



Sample: Lane 1: Mouse Lung Lysates Lane 2: Mouse Kidney Lysates Lane 3: Rat Lung Lysates Lane 4: Rat Kidney Lysates Primary: Anti-Phospho-TIE2 (Tyr1108)(bs-12262R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 124kDa Observed band size: 160kDa



Paraformaldehyde-fixed, paraffin embedded (Human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Phospho-TIE2 (Tyr1108)) Polyclonal Antibody, Unconjugated (bs-12262R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.