

bsm-60167R**[Primary Antibody]****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

Ubiquityl-Histone H2B (Lys120) Recombinant Rabbit mAb**DATASHEET**


Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:200-1000) IHC-F (1:200-1000) IF (1:200-1000) Reactivity: Human, Mouse, Rat Predicted MW.: 14 kDa Subcellular Location: Nucleus
Clonality: Recombinant	CloneNo.: U2A2	
GeneID: 3018	SWISS: P33778	
Target: Ubiquityl-Histone H2B (Lys120)		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H2B family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq, Jul 2008].		

VALIDATION IMAGES

Blocking buffer: 5% NFDM/TBST Primary ab dilution: 1:2000 Primary ab incubation condition: 2 hours at room temperature Secondary ab: Goat Anti-Rabbit IgG H&L (HRP) Lysate: HeLa, N2a, BRL, Mouse liver, Mouse kidney Protein loading quantity: 20 µg Exposure time: 30 Predicted MW: 23Da Observed MW: 23Da



Tissue: Rat liver Section type: Formalin fixed & Paraffin -embedded section Retrieval method: High temperature and high pressure Retrieval buffer: Tris/EDTA buffer, pH 9.0 Primary ab dilution: 1:1000 Primary ab incubation condition: 1 hour at room temperature Secondary ab: SP Kit(Rabbit) (sp-0023) Counter stain: Hematoxylin (Blue) Comment: Color brown is the positive signal for bsm-60167R



Tissue: Mouse liver Section type: Formalin fixed & Paraffin -embedded section Retrieval method: High temperature and high pressure Retrieval buffer: Tris/EDTA buffer, pH 9.0 Primary ab dilution: 1:1000 Primary ab incubation condition: 1 hour at room temperature Secondary ab: SP Kit(Rabbit) (sp-0023) Counter stain: Hematoxylin (Blue) Comment: Color brown is the positive signal for bsm-60167R

SELECTED CITATIONS

- **[IF=6.2]** Wenjun Chen. et al. MdLRR-RLK1-MdATG3 module enhances the resistance of apples to abiotic stress via autophagy. PLANT J. 2024 Dec; WB ;Escherichia coli. 39671299