
OVOL2 Rabbit pAb

Catalog Number: bs-12274R

Target Protein: OVOL2

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Mouse, Rat (predicted:Human, Pig, Cow)

Predicted MW: 30 kDa

Entrez Gene: 58495

Swiss Prot: Q9BRP0

Source: KLH conjugated synthetic peptide derived from human OVOL2: 75-180/275.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: The Ovo family of zinc-finger transcription factors encode evolutionarily conserved genes including those from *Caenorhabditis elegans*, *Drosophila melanogaster*, mouse and human. Members of the Ovo family include Ovol1 and Ovol2. Ovol1 acts as a transcriptional repressor by interacting with key developmental signaling pathways such as Wnt and TGF- β /BMP. Specifically, Ovol1 represses c-Myc and Id2 genes and establishes a balance between proliferation and differentiation of progenitor cells. Deletion of Ovol1 in mice leads to germ cell degeneration and defective sperm production in adult males. Ovol1 has also been shown to repress itself as well as Ovol2, which is thought to regulate neural development and vascular angiogenesis during embryogenesis.

VALIDATION IMAGES



Sample: Lane 1: Heart (Mouse) Lysate at 40 ug Primary: Anti-OVOL2 (bs-12274R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 37 kD Observed band size: 35 kD



Paraformaldehyde-fixed, paraffin embedded (rat liver tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (OVOL2) Polyclonal Antibody, Unconjugated (bs-12274R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.