


bs-60044R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

Histone H2B (Acetyl K116) Rabbit pAb**DATASHEET****Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 3018**SWISS:** P33778**Target:** Histone H2B (Acetyl K116)**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.**Background:** Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H2B family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq, Jul 2008].**Applications:** WB (1:500-2000)**ChIP** (4 ug/5x10⁶ cells)**Reactivity:** Human (predicted: Mouse, Rat)**Predicted MW.:** 14 kDa**Subcellular Location:** Nucleus**VALIDATION IMAGES**

Blocking buffer: 5% NFDM/TBST Primary ab dilution: 1:2000 Primary ab incubation condition: 2 hours at room temperature
Secondary ab: Goat Anti-Rabbit IgG H&L (HRP)
Lysate: (-) HeLa, (+) HeLa+Sodium butyrate (30mM, 4hr) Protein loading quantity: 20 µg
Exposure time: 60 s Predicted MW: 14 kDa
Observed MW: 14 kDa



Cell type: HeLa+SBA (5mM, 24h) Cross-linking conditions: No cross-linking Amount of chromatin per IP: 5 × 10⁶ cells Amount of ab per IP: 4 µg, 12 µg Beads type and amount per IP: 50 µl of Protein A/G MagBeads Comment: The ChIP was performed with 1 µg of normal rabbit IgG as a negative control. Real time quantitative PCR was performed on immunoprecipitated DNA using primers specific for the human RAB20, GAPDH CDS region, RPL30, LDHA, FOXO3a-promoter and FOXO3a-downstream. Data are presented as enrichment of each sample relative to total amount of input chromatin at each amplicon.