

**bsm-60063M****[ Primary Antibody ]****Histone H2B (Acetyl K15) Mouse mAb****Bioss**  
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —**

<b>Host:</b> Mouse	<b>Isotype:</b> IgG	<b>Applications:</b> WB (1:500-2000)  <b>Reactivity:</b> Human (predicted: Mouse, Rat)  <b>Predicted MW.:</b> 14 kDa  <b>Subcellular Location:</b> Nucleus
<b>Clonality:</b> Monoclonal	<b>CloneNo.:</b> H2A4	
<b>GeneID:</b> 3018	<b>SWISS:</b> P33778	
<b>Target:</b> Histone H2B (Acetyl K15)		
<b>Purification:</b> affinity purified by Protein G		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H2B family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq, Jul 2008].		

**— VALIDATION IMAGES —**

Blocking buffer: 5% NFDM/TBST Primary ab  
dilution: 1:2000 Primary ab incubation  
condition: 2 hours at room temperature  
Secondary ab: Goat Anti-Mouse IgG H&L (HRP)  
Lysate: (-) HeLa, (+) HeLa+Sodium butyrate  
(30mM, 4hr)+Trichostatin A (2μM, 4hr) Protein  
loading quantity: 20 μg Exposure time: 60 s  
Predicted MW: 14 kDa Observed MW: 14 kDa