

bsm-60047M**[Primary Antibody]****BioSS**
ANTIBODIES

www.bioss.com.cn



sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

Histone H2B (Acetyl K20) Mouse mAb**DATASHEET**

Host: Mouse	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:50-100) IF (1:50-100) Reactivity: Human (predicted: Mouse, Rat) Predicted MW.: 14 kDa Subcellular Location: Nucleus
Clonality: Monoclonal	CloneNo.: H2F5	
GeneID: 3018	SWISS: P33778	
Target: Histone H2B (Acetyl K20)		
Purification: affinity purified by Protein G		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H2B family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq, Jul 2008].		

VALIDATION IMAGES

Cell type: HeLa + Serum starvation (12 h) + SBA (5 mM, 24 h) Cross-linking conditions: No cross-linking Amount of chromatin per IP: 5×10^6 cells Amount of Ab per IP: 4 μ g Beads type and amount per IP: 50 μ L of Protein A/G MagBeads Comment: The ChIP was performed with 1 μ g of normal rabbit IgG as a negative control. Real time quantitative PCR was performed on immunoprecipitated DNA using primers specific for the human GAPDH CDS region, RPL30 Exon 3, FOXO3a-promoter and FOXO3a-downstream. Data are presented as enrichment of each sample relative to total amount of input chromatin at each amplicon.

Blocking buffer: 5% NFDM/TBST Primary ab dilution: 1:2000 Primary ab incubation condition: 2 hours at room temperature Secondary ab: Goat Anti-Mouse IgG H&L (HRP) Lysate: (-) HeLa, (+) HeLa+Sodium butyrate (30mM, 4hr) Protein loading quantity: 20 μ g Exposure time: 60 s Predicted MW: 14 kDa Observed MW: 14 kDa