
Histone H2B (Acetyl K23) Mouse mAb

Catalog Number: bsm-60061M

Target Protein: Histone H2B (Acetyl K23)

Concentration: 1mg/ml

Form: Liquid

Host: Mouse

Clonality: Monoclonal

Clone No.: H2B9

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:50-100), IF (1:50-100)

Reactivity: Human, Mouse (predicted:Rat, Rice)

Predicted MW: 14 kDa

Entrez Gene: 3018

Swiss Prot: P33778


Purification: affinity purified by Protein G

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H2B family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq, Jul 2008].

VALIDATION IMAGES



Blocking buffer: 5% NFDM/TBST Primary ab dilution: 1:2000 Primary ab incubation condition: 2 hours at room temperature Secondary ab: Goat Anti-Mouse IgG H&L (HRP) Lysate: (-) HeLa, (+) HeLa+Trichostatin A (2 μ M, 4hr) Protein loading quantity: 20 μ g Exposure time: 60 s Predicted MW: 14 kDa Observed MW: 14 kDa



Blocking buffer: 5% NFDM/TBST Primary ab dilution: 1:1000 Primary ab incubation condition: 2 hours at room temperature Secondary ab: Goat Anti-Mouse IgG H&L (HRP) Lysate: N2a, Mouse kidney, Rice Protein loading quantity: 20 μ g Exposure time: 60 s Predicted MW: 14 kDa Observed MW: 14 kDa