

bs-60039R**[Primary Antibody]****Histone H2B (Acetyl K24) Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		Reactivity: Human (predicted: Mouse, Rat)
GeneID: 3018	SWISS: P33778	
Target: Histone H2B (Acetyl K24)		Predicted MW.: 14 kDa
Purification: affinity purified by Protein A		Subcellular Location: Nucleus
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H2B family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq, Jul 2008].		

— VALIDATION IMAGES —

Blocking buffer: 5% NFDM/TBST Primary ab
 dilution: 1:2000 Primary ab incubation
 condition: 2 hours at room temperature
 Secondary ab: Goat Anti-Rabbit IgG H&L (HRP)
 Lysate: (-) HeLa, (+) HeLa+Sodium butyrate
 (30mM, 4hr) Protein loading quantity: 20 µg
 Exposure time: 60 s Predicted MW: 14 kDa
 Observed MW: 14 kDa