

phospho-delta 1 Catenin (Ser268) Rabbit pAb

Catalog Number: bs-12989R

Target Protein: phospho-delta 1 Catenin (Ser268)

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human, Mouse (predicted:Rat, Pig, Dog)

Predicted MW: 108 kDa

Entrez Gene: 1500

Swiss Prot: O60716

Source: KLH conjugated synthesised phosphopeptide derived from human delta 1 Catenin around the phosphorylation site of Ser268: GG(p-S)SV.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Binds to and inhibits the transcriptional repressor ZBTB33, which may lead to activation of target genes of the Wnt signaling pathway (By similarity). May associate with and regulate the cell adhesion properties of both C- and E-cadherins. Implicated both in cell transformation by SRC and in ligand-induced receptor signaling through the EGF, PDGF, CSF-1 and ERBB2 receptors. Promotes GLIS2 C-terminal cleavage.

VALIDATION IMAGES



Sample: NIH/3T3(Mouse) Cell Lysate at 30 ug A431(Human) Cell Lysate at 30 ug Primary: Anti- phospho-delta 1 Catenin (Ser268) (bs-12989R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 108 kD Observed band size: 110 kD



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-delta 1 Catenin (Ser268)) Polyclonal Antibody, Unconjugated (bs-12989R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.