

bs-12307R**[Primary Antibody]****ENPEP Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000) Reactivity: (predicted: Human, Mouse, Rat, Sheep, Cow, Dog) Predicted MW.: 109 kDa Subcellular Location: Cell membrane
Clonality: Polyclonal		
GeneID: 2028	SWISS: Q07075	
Target: ENPEP		
Immunogen: KLH conjugated synthetic peptide derived from human Aminopeptidase A: 766-830/957. < Extracellular >		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Aminopeptidase A, also designated APA, gp160 human kidney differentiation antigen, glutamyl aminopeptidase, or enpep, is a differentiation-related kidney glycoprotein. As a cell surface, zinc-dependent metalloprotease, Aminopeptidase A specifically cleaves amino-terminal acidic residues from peptide substrates such as Angiotensin II. APA is expressed on the surface of epithelial cells of the glomerulus and proximal tubule cells of the human nephron, where it may mediate the constitutive trafficking of Glut4-containing vesicles. These Glut4-containing vesicles are tissue-specific secretory-like microsomal structures that mediate insulin-dependent translocation of GLUT4 to the cell surface in fat and muscle cells. Mutations in the gp160/APA gene, including loss of protein expression or enzymatic activity, occur in 20% of primary clear cell renal carcinomas.		