

**bs-12854R****[ Primary Antibody ]****phospho-Beta catenin (Ser37) Rabbit pAb****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —****Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 1499**SWISS:** P35222**Target:** Beta catenin (Ser37)**Immunogen:** KLH conjugated synthesised phosphopeptide derived from human beta Catenin around the phosphorylation site of Ser37: IH(p-S)GA.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

**Background:** The protein encoded by this gene is part of a complex of proteins that constitute adherens junctions (AJs). AJs are necessary for the creation and maintenance of epithelial cell layers by regulating cell growth and adhesion between cells. The encoded protein also anchors the actin cytoskeleton and may be responsible for transmitting the contact inhibition signal that causes cells to stop dividing once the epithelial sheet is complete. Finally, this protein binds to the product of the APC gene, which is mutated in adenomatous polyposis of the colon. Mutations in this gene are a cause of colorectal cancer (CRC), pilomatrixoma (PTR), medulloblastoma (MDB), and ovarian cancer. Three transcript variants encoding the same protein have been found for this gene.[provided by RefSeq, Oct 2009].**Applications:** WB (1:500-2000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**Flow-Cyt** (2ug/Test)**Reactivity:** Human, Mouse, Rat  
(predicted: Rabbit, Pig, Chicken)**Predicted**  
**MW.:** 86 kDa**Subcellular** Cell membrane ,Cytoplasm  
**Location:** ,Nucleus**— VALIDATION IMAGES —**

Sample: Cerebrum (Mouse) Lysate at 40 ug  
Cerebrum (Rat) Lysate at 40 ug  
Primary: Anti-phospho-beta Catenin (Ser37) (bs-12854R) at 1/1000 dilution  
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution  
Predicted band size: 92 kD  
Observed band size: 92 kD



Paraformaldehyde-fixed, paraffin embedded (Rat colon); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (phospho-beta Catenin (Ser37)) Polyclonal Antibody, Unconjugated (bs-12854R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control: 293T. Primary Antibody (green line): Rabbit Anti-phospho-beta Catenin (Ser37) antibody (bs-12854R) Dilution: 2µg / 10<sup>6</sup> cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody : Goat anti-rabbit IgG-FITC Dilution: 1µg / test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 0.1% PBST for 20 min at room temperature. The cells were then incubated in 5% BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

**— SELECTED CITATIONS —**

- **[IF=5.59]** Otsuka, Shotaro. et al. Preconditioning Exercise in Rats Attenuates Early Brain Injury Resulting from

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Subarachnoid Hemorrhage by Reducing Oxidative Stress, Inflammation, and Neuronal Apoptosis. 2021 Aug 09 WB ;Rat. 34368932

- **[IF=3]** Gongyan Liu. et al. Hair Follicle Development of Rex Rabbits Is Regulated Seasonally by Wnt10b/ $\beta$ -Catenin, TGF $\beta$ -BMP, IGF1, and EGF Signaling Pathways. ANIMALS. 2023 Jan;13(23):3742 WB ;Rabbit. 38067094