

bs-25287R**[Primary Antibody]****TMEM30A Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —


Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		IHC-P (1:100-500)
GeneID: 55754	SWISS: Q9NV96	IHC-F (1:400-800)
Target: TMEM30A		IF (1:100-500)
Immunogen: KLH conjugated synthetic peptide derived from human TMEM30A: 300-361/361. < Cytoplasmic >		Reactivity: Mouse, Rat (predicted: Human, Pig, Sheep, Cow, Horse)
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		Predicted MW.: 41 kDa
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: Cell membrane
Background: Enables aminophospholipid flippase activity. Involved in several processes, including phospholipid transport; positive regulation of transport; and xenobiotic transmembrane transport. Located in endoplasmic reticulum and plasma membrane. Part of phospholipid-translocating ATPase complex. [provided by Alliance of Genome Resources, Nov 2021]		

— VALIDATION IMAGES —

Sample: Lane 1: Rat Liver tissue lysates Lane 2: Rat Kidney tissue lysates Primary: Anti-TMEM30A (bs-25287R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 41 kDa Observed band size: 40 kDa



Paraformaldehyde-fixed, paraffin embedded (mouse kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (TMEM30A) Polyclonal Antibody, Unconjugated (bs-25287R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (TMEM30A) Polyclonal Antibody, Unconjugated (bs-25287R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.