

bs-12876R**[Primary Antibody]****BOB1 Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: ELISA (1:5000-10000)
Clonality: Polyclonal		Reactivity: (predicted: Human, Mouse, Rat, Rabbit, Pig, Sheep, Cow, Chicken, Dog, Horse)
GeneID: 5450	SWISS: Q16633	Predicted MW.: 27 kDa
Target: BOB1		Subcellular Location: Nucleus
Immunogen: KLH conjugated synthetic peptide derived from human BOB1: 1-100/256.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: POU domain proteins contain a bipartite DNA-binding domain divided by a flexible linker that enables them to adopt various monomer configurations on DNA. The versatility of POU protein operation is additionally conferred at the dimerization level. The POU dimer from the OCT1 gene formed on the palindromic OCT factor recognition element, or PORE (ATTTGAAATGCAAAT), could recruit the transcriptional coactivator OBF1. Studies of tissue-specific expression of immunoglobulin promoters demonstrate the importance of an octamer, ATTTGCAT, and the proteins that bind to it. This is a regulatory element important for tissue- and cell-specific transcription as well as for transcription of a number of housekeeping genes. Oct-1 encodes one protein, NF-A1, which is found in nuclear extracts from all cell types and thus is not specific to lymphoid cells as is the protein NF-A2, which is encoded by Oct-2. A novel protein designated Bob 1 (B cell Oct binding protein 1), alternatively called OBF-1, specifically interacts with Oct-1 and Oct-2, enhancing their transcriptional efficacy. Bob 1 is expressed at highest levels in spleen and peripheral blood leukocytes and represents an Oct co-factor capable of conferring cell-specific activation of Oct-1 and Oct-2. Although having no intrinsic capacity for DNA binding, Bob 1 associates tightly with the octamer motif in the presence of Oct-1 and/or Oct-2. The gene which encodes Bob 1 maps to human chromosome 11q23.1.		