
A4GALT Rabbit pAb

Catalog Number: bs-25228R

Target Protein: A4GALT

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000)

Reactivity: Human, Mouse (predicted:Rat, Sheep, Cow, Dog)

Predicted MW: 40 kDa

Entrez Gene: 53947

Swiss Prot: Q9NPC4

Source: KLH conjugated synthetic peptide derived from human A4GALT: 31-130/353.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Necessary for the biosynthesis of the Pk antigen of blood histogroup P. Catalyzes the transfer of galactose to lactosylceramide and galactosylceramide. Necessary for the synthesis of the receptor for bacterial verotoxins. Expression of CD77, also called Gb3, sensitizes a cell to verotoxins, causing cellular injury that can lead to disease. Therefore, the complex regulation of CD77 biosynthesis and the activity of the enzymes involved, such as CD77 synthase, can be studied by compared gene expression between toxin-sensitive and insensitive tissues and cell lines. The highest tissue expression of CD77 synthase occurs in the kidney, mesenteric lymph node, spleen, and brain. Burkitt leukemia cells express very high levels of CD77 as well as CD77 synthase, and are sensitive to verotoxin induced apoptosis. These megakaryoblasts then never mature, leading to the arrest of platelet generation in the bone marrow, which may cause thrombocytopenia, a symptom associated with various hemorrhagic conditions.

VALIDATION IMAGES



Sample: Lane 1: Mouse Kidney tissue lysates Lane 2: Mouse Heart tissue lysates Lane 3: Mouse Testis tissue lysates Lane 4: Human HeLa cell lysates Primary: Anti-A4GALT (bs-25228R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 40 kDa Observed band size: 50 kDa