

bs-13008R**[Primary Antibody]****DLST Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

DATASHEET


Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ELISA (1:5000-10000) Reactivity: Human, Mouse, Rat (predicted: Rabbit, Pig, Sheep, Cow, Dog, Horse) Predicted MW.: 41 kDa Subcellular Location: Cytoplasm
Clonality: Polyclonal		
GeneID: 1743	SWISS: P36957	
Target: DLST		
Immunogen: KLH conjugated synthetic peptide derived from human DLST: 201-300/453.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: The 2-oxoglutarate dehydrogenase complex catalyzes the overall conversion of 2-oxoglutarate to succinyl-CoA and CO ₂ . The complex contains multiple copies of three enzymatic components: 2-oxoglutarate dehydrogenase (E1), dihydrolipoamide succinyltransferase (E2) and lipoamide dehydrogenase (E3). DLST (dihydrolipoyllysine-residue succinyltransferase component of 2-oxoglutarate dehydrogenase complex, mitochondrial), also known as DLTs or 2-oxoglutarate dehydrogenase complex component E2, is a 453 amino acid protein belonging to the 2-oxoacid dehydrogenase family. DLST covalently binds one lipoyl cofactor and participates in L-lysine degradation via the saccharopine pathway. Localized to the mitochondrion, DLST forms a 24-polypeptide structural core with octahedral symmetry. The gene encoding DLST maps to human chromosome 14q24.3 and mouse chromosome 12 D2.		

VALIDATION IMAGES

Sample: Lane 1: Mouse Liver tissue lysates Lane 2: Mouse Muscle tissue lysates Lane 3: Mouse Testis tissue lysates Lane 4: Rat Heart tissue lysates Lane 5: Rat Muscle tissue lysates Primary: Anti-DLST (bs-13008R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 41 kDa Observed band size: 46 kDa



Paraformaldehyde-fixed, paraffin embedded (mouse skeletal muscle); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (DLST) Polyclonal Antibody, Unconjugated (bs-13008R) at 1:100 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human skeletal muscle); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (DLST) Polyclonal Antibody, Unconjugated (bs-13008R) at 1:100 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

SELECTED CITATIONS

- **[IF=3.5]** Xiaoxuan Zhao. et al. Demystifying the Landscape of Endometrial Immune Microenvironment in Luteal-Phase from Cuprotosis: Implications for the Mechanism and Treatment of RPL. GENE. 2024 Jan;;148191 IF ;Mouse. 38253297

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.