

bs-3389R**[Primary Antibody]****phospho-RPS6 (Ser240 + Ser244) Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —


| | |
|---|---|
| <p>Host: Rabbit</p> <p>Clonality: Polyclonal</p> <p>GeneID: 6194</p> <p>Target: RPS6 (Ser240 + Ser244)</p> <p>Immunogen: KLH conjugated Synthesised phosphopeptide derived from human RPS6 around the phosphorylation site of Ser240/244: RA(p-S)TSK(p-S)ES.</p> <p>Purification: affinity purified by Protein A</p> <p>Concentration: 1mg/ml</p> <p>Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p>Background: Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a cytoplasmic ribosomal protein that is a component of the 40S subunit. The protein belongs to the S6E family of ribosomal proteins. It is the major substrate of protein kinases in the ribosome, with subsets of five C-terminal serine residues phosphorylated by different protein kinases. Phosphorylation is induced by a wide range of stimuli, including growth factors, tumor-promoting agents, and mitogens. Dephosphorylation occurs at growth arrest. The protein may contribute to the control of cell growth and proliferation through the selective translation of particular classes of mRNA. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. [provided by RefSeq, Jul 2008]</p> | <p>Isotype: IgG</p> <p>SWISS: P62753</p> <p>Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)</p> <p>Reactivity: Human, Mouse, Rat</p> <p>Predicted MW.: 29 kDa</p> <p>Subcellular Location: Cell membrane ,Cytoplasm ,Nucleus</p> |
|---|---|

— VALIDATION IMAGES —

NIH/3T3 (M) cells were treated with or without FBS 20% for 30 min, 25 µg total protein per lane of cell lysates (see on figure) probed with Phospho-RPS6 (Ser240 + Ser244) polyclonal antibody, unconjugated (bs-3389R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Human Tonsil; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with phospho-RPS6 (Ser240 + Ser244) Polyclonal Antibody, Unconjugated (bs-3389R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Breast Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with phospho-RPS6 (Ser240 + Ser244) Polyclonal Antibody, Unconjugated (bs-3389R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.