bs-10876R

[Primary Antibody]

Cystatin-C Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET			400-901-9800
Host: Rat	bit Is	otype: IgG	Applications: WB (1:500-2000)
Clonality: Pol	yclonal		IHC-P (1:100-500)
GenelD: 147	1 S	WISS: P01034	IF (1:100-500)
Target: Cys	tatin-C		Reactivity: Human, Mouse, Rat
Purification: affi	nity purified by Protein A		, , , , , , , , , ,
Concentration: 1m	g/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.			Predicted MW.: ^{13 kDa} Subcellular
Background: The mu cys new fan typ are hur pro cor pse enc pro anc this Exp sev lesi Ref	ground: The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins(stefins), type 2 cystatins and the kininogens. The type 2 cystatin proteins are a class of cysteine proteinase inhibitors found in a variety of human fluids and secretions, where they appear to provide protective functions. The cystatin locus on chromosome 20 contains the majority of the type 2 cystatin genes and pseudogenes. This gene is located in the cystatin locus and encodes the most abundant extracellular inhibitor of cysteine proteases, which is found in high concentrations in biological fluids and is expressed in virtually all organs of the body. A mutation in this gene has been associated with amyloid angiopathy. Expression of this protein in vascular wall smooth muscle cells is severely reduced in both atherosclerotic and aneurysmal aortic lesions, establishing its role in vascular disease. [provided by RefSeq].		Location: Secreted

– VALIDATION IMAGES



Sample: Testis (Mouse) Lysate at 40 ug Cerebrum (Mouse) Lysate at 40 ug Primary: Anti-Cystatin-C (bs-10876R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 13 kD Observed band size: 13 kD



Sample: MDA-MB-231(Human) Cell Lysate at 30 ug Primary: Anti-Cystatin-C (bs-10876R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 13 kD Observed band size: 13 kD



Paraformaldehyde-fixed, paraffin embedded (Human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Cystatin-C) Polyclonal Antibody, Unconjugated (bs-10876R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CST3) Polyclonal Antibody, Unconjugated (bs-10876R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.

- SELECTED CITATIONS -

- [IF=6.1] Qingyu Ding. et al. N-acetylcysteine alleviates oxidative stress and apoptosis and prevents skeletal muscle atrophy in type 1 diabetes mellitus through the NRF2/HO-1 pathway. LIFE SCI. 2023 Sep;329:121975 WB ;Dog. 37495077
- [IF=4.5] Gan Rao. et al. Arsenic and polystyrene-nano plastics co-exposure induced testicular toxicity: Triggers oxidative stress and promotes apoptosis and inflammation in mice. ENVIRON TOXICOL. 2023 Sep;: WB ;MOUSE. 37705229
- [IF=3.8] Qiuxiang Cai. et al. Pasteurella multocida causes liver injury in ducks by mediating inflammatory, apoptotic and autophagic pathways. MICROB PATHOGENESIS. 2023 Sep;:106336 WB,IHC,IF ;Duck. 37683832