

bs-23388R**[Primary Antibody]****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

PER2 Rabbit pAb**— DATASHEET —**

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal		IHC-F (1:100-500)
GeneID: 8864	SWISS: O15055	IF (1:100-500)
Target: PER2		Reactivity: Mouse, Rat (predicted: Human)
Immunogen: KLH conjugated synthetic peptide derived from human PER2: 1001-1100/1255.		
Purification: affinity purified by Protein A		Predicted MW.: 138 kDa
Concentration: 1mg/ml		Subcellular Location: Cytoplasm ,Nucleus
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: PER2, a mammalian homologue of the Drosophila period gene, shares a 40% homology with PER1 including the protein dimerization PAS domain. PER2 is a circadian regulator that may act as a transcription factor. It behaves as a negative element in circadian transcriptional loop. PER2 does not appear to bind DNA, suggesting indirect transcriptional inhibition. Expression oscillates with a 24 hour rhythm in the suprachiasmatic nucleus (SCN) and the whole eyes. Oscillations are maintained under constant darkness and are responsive to changes of the light/dark cycles. There is a 4 hour time delay between PER1 and PER2 oscillations. The expression rhythms appear to originate from retina.		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (Mouse pancreas); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PER2/Period circadian protein 2) Polyclonal Antibody, Unconjugated (bs-23388R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PER2/Period circadian protein 2) Polyclonal Antibody, Unconjugated (bs-23388R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.