bsm-51028M

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

MMP14 Mouse mAb

DATASHEET -

Host: Mouse Isotype: IgG CloneNo.: 3A6 Clonality: Monoclonal GeneID: 4323 **SWISS:** P50281

Target: MMP14

Purification: affinity purified by Protein AGL

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: MMP14 may be an activator of pro gelatinase A and is expressed in

fibroblast cells during both wound healing and human cancer progression. MMP14 is expressed in very low levels and may require stimulation with concanavolin A or the phorbol ester TPA

to stimulate production of MMP14.

Matrix metalloproteinase-14 precursor is endopeptidase that degrades various components of the extracellular matrix, such as collagen. Activates progelatinase A. Essential for pericellular collagenolysis and modeling of skeletal and extraskeletal connective tissues during development. [Catalytic activity] Endopeptidase activity. Activates progelatinase A by cleavage of the propeptide at 37-Asn-|-Leu-38. Other bonds hydrolyzed include 35-Gly-|-Ile-36 in the propertide of collagenase 3, and 341-Asn-|-Phe-342, 441-Asp-|-Leu-442 and 354-Gln-|-Thr-355 in the aggrecan interglobular domain. Highly expressed in placenta, kidney, heart, lung, embryonic skeletal and periskeletal tissues. Belongs to the

peptidase M10A family.

Applications: WB (1:500-2000)

ICC/IF (1:50-200)

Reactivity: Human, Mouse

(predicted: Rat, Rabbit, Pig,

Cow, Bovine)

Predicted 65 kDa

MW.:

Subcellular Location: Cell membrane