bs-12925R

[Primary Antibody]

Bioss ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

COX6c Rabbit pAb

GeneID: 1345 **SWISS:** P09669

Target: COX6c

Immunogen: KLH conjugated synthetic peptide derived from human

COX6c/Cytochrome C Oxidase subunit VIc: 21-75/75.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: This protein is one of the nuclear-coded polypeptide chains of

cytochrome c oxidase, the terminal oxidase in mitochondrial

electron transport.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:100-500)

Reactivity: Human, Rat


(predicted: Mouse, Rabbit, Pig, Sheep, Cow, Dog,

Horse)


Predicted 9 kDa

Subcellular Location: Cell membrane ,Cytoplasm

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (COX6c) Polyclonal Antibody, Unconjugated (bs-12925R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (COX6c) Polyclonal Antibody, Unconjugated (bs-12925R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.