## bs-49183R

# [ Primary Antibody ]

# Bioss

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

# **EAAT2** Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**GenelD:** 20511 **SWISS:** P43006

Target: EAAT2

Immunogen: Recombinant mouse EAAT2 protein: 460-572/572.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** This gene encodes a member of a family of solute transporter

proteins. The membrane-bound protein is the principal transporter that clears the excitatory neurotransmitter glutamate from the extracellular space at synapses in the central nervous system. Glutamate clearance is necessary for proper synaptic activation and to prevent neuronal damage from excessive activation of glutamate receptors. Mutations in and decreased expression of this protein are associated with amyotrophic lateral sclerosis.

Alternatively spliced transcript variants of this gene have been

identified. [provided by RefSeq, Sep 2010]

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)


**Reactivity:** Human, Mouse

(predicted: Rat)


Predicted MW.: 62 kDa

Subcellular Location: Cell membrane

### VALIDATION IMAGES -




Sample: Lane 1: Recombinant mouse EAAT2 protein, GST & His Primary: Anti-EAAT2 (bs-49183R) at 1/500 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 62 kDa Observed band size: 36 kDa




Paraformaldehyde-fixed, paraffin embedded (human cerebellum); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (EAAT2) Polyclonal Antibody, Unconjugated (bs-49183R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (EAAT2) Polyclonal Antibody, Unconjugated (bs-49183R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



Paraformaldehyde-fixed, paraffin embedded Mouse Cerebellum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with EAAT2 Polyclonal Antibody, Unconjugated (bs-49183R) at 1:200 overnight at 4°C. Followed by conjugated Goat



Paraformaldehyde-fixed, paraffin embedded Human Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with EAAT2 Polyclonal Antibody, Unconjugated (bs-49183R) at 1:200 overnight at 4°C. Followed by conjugated Goat Anti-Rabbit IgG antibody (Red, bs-0295G-BF594), Anti-Rabbit IgG antibody (Red, bs-0295G-BF594), DAPI (blue, C02-04002) was used to stain the cell DAPI (blue, C02-04002) was used to stain the cell nuclei. nuclei.