

bs-0094R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

AACT Rabbit pAb**— DATASHEET —**


Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		IHC-P (1:100-500)
GeneID: 12	SWISS: P01011	IHC-F (1:100-500)
Target: AACT		IF (1:100-500)
Immunogen: KLH conjugated synthetic peptide derived from human AACT: 121-320/423.		Reactivity: Human, Mouse, Rat
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		Predicted MW.: 45 kDa
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: Secreted
Background: Alpha 1-Antichymotrypsin, a member of the serine proteinase inhibitor (serpin) family, inhibits neutrophil proteinase cathepsin G and mast cell chymases, and protects the lower respiratory tract from damage by proteolytic enzymes. It contains a reactive centre loop, which interacts with cognate proteinases, resulting in loop cleavage and a major conformational change. Recently, alpha 1-antichymotrypsin has been identified as a major constituent of the neurofibrillary plaques associated with Alzheimers disease, and in vitro studies have shown that it enhances the rate of amyloid-fibril formation. These observations and recent genetic evidence suggest that alpha 1-antichymotrypsin is important in the pathogenesis of Alzheimers disease.		

— VALIDATION IMAGES —


Sample: Liver (Mouse) Lysate at 40 ug Primary: Anti-Alpha-1-Antichymotrypsin (bs-0094R) at 1/500 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 45 kD Observed band size: 47 kD



Sample: Lane 1: Mouse Liver tissue lysates Lane 2: Rat Liver tissue lysates Primary: Anti-Alpha-1-Antichymotrypsin (bs-0094R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 45 kDa Observed band size: 45 kDa



Tissue/cell: Mouse liver tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-Alpha-1-Antichymotrypsin Polyclonal Antibody, Unconjugated (bs-0094R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody (SP-0023) and DAB (C-0010) staining



Tissue/cell: Rat brain tissue; 4%

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti- Alpha-1-Antichymotrypsin Polyclonal Antibody, Unconjugated (bs-0094R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

— SELECTED CITATIONS —

- **[IF=4.2]** Jin Rao. et al. Integration of Microarray Data and Single-Cell Sequencing Analysis to Explore Key Genes Associated with Macrophage Infiltration in Heart Failure. J INFLAMM RES. 2024 Dec 19 WB ;Mouse. 39717663
- **[IF=3.13]** Wang, Robert YL, et al. "Proteome Demonstration of Alpha-1-Acid Glycoprotein and Alpha-1-Antichymotrypsin." Pediatric Infectious Disease Journal (2014). WB ;="Human". 25170552
- **[IF=3.3]** Yan Zhang. et al. Proteomic analysis of multiple organ dysfunction induced by rhabdomyolysis. J PROTEOMICS. 2024 Apr;298:105138 IHC,WB ;Rat. 38403185