

---

## JAK1 Recombinant Rabbit mAb

Catalog Number: bsm-54138R

Target Protein: JAK1

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Recombinant

Clone No.: 2B12

Isotype: IgG

Applications: WB (1:200-1000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human, Mouse, Rat

Predicted MW: 133 kDa

Subcellular Cytoplasm

Locations:

Entrez Gene: 3716

Swiss Prot: P23458

Purification: affinity purified by Protein A


Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


**Background:** Janus kinase 1 (JAK1) is a member of a new class of non-receptor protein-tyrosine kinases (PTK) characterized by the presence of a second phosphotransferase-related domain immediately N-terminal to the PTK domain. The second phosphotransferase domain bears all the hallmarks of a protein kinase, although its structure differs significantly from that of the PTK and threonine/serine kinase family members. JAK1 is a large, widely expressed membrane-associated phosphoprotein. It is involved in the interferon-alpha/beta and -gamma signal transduction pathways. The reciprocal interdependence between JAK1 and TYK2 activities in the interferon-alpha pathway, and between JAK1 and JAK2 in the interferon-gamma pathway, may reflect a requirement for these kinases in the correct assembly of interferon receptor complexes. These kinases couple cytokine ligand binding to tyrosine phosphorylation of various known signaling proteins and a unique family of transcription factors termed the signal transducers and activators of transcription, or STATs.

### VALIDATION IMAGES


---




25 ug total protein per lane of various lysates (see on figure) probed with JAK1 monoclonal antibody, unconjugated (bsm-54138R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.




Paraformaldehyde-fixed, paraffin embedded Mouse Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with JAK1 Monoclonal Antibody, Unconjugated(bsm-54138R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.




Paraformaldehyde-fixed, paraffin embedded Rat Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with JAK1 Monoclonal Antibody, Unconjugated(bsm-54138R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with JAK1 Monoclonal Antibody, Unconjugated(bsm-54138R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Mouse Spleen; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with JAK1 Monoclonal Antibody, Unconjugated(bsm-54138R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Mouse Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with JAK1 Monoclonal Antibody, Unconjugated(bsm-54138R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.

## PRODUCT SPECIFIC PUBLICATIONS

[IF=17.4] Hui Yan. et al. Elaboration a ROS-responsive darutigenol prodrug nanoassemblies for inflammatory arthritis treatment. NANO TODAY. 2024 Apr;55:102220 WB ; Mouse . 10.1016/j.nantod.2024.102220

[IF=7.5] Junchao Wu. et al. JAK1/JAK2 degraders based on PROTAC for topical treatment of atopic dermatitis. BIOMED PHARMACOTHER. 2024 Feb;171:116167 WB,IHC ; Mouse . 38262152