

**bs-0583R****[ Primary Antibody ]****c-Abl Rabbit pAb****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit <b>Clonality:</b> Polyclonal <b>GeneID:</b> 25 <b>Target:</b> c-Abl <b>Immunogen:</b> KLH conjugated synthetic peptide derived from human c-Abl: 401-500/1149. <b>Purification:</b> affinity purified by Protein A <b>Concentration:</b> 1mg/ml <b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. <b>Background:</b> The ABL1 protooncogene encodes a cytoplasmic and nuclear protein tyrosine kinase that has been implicated in processes of cell differentiation, cell division, cell adhesion, and stress response. Activity of c-Abl protein is negatively regulated by its SH3 domain, and deletion of the SH3 domain turns ABL1 into an oncogene. The t(9;22) translocation results in the head-to-tail fusion of the BCR (MIM:151410) and ABL1 genes present in many cases of chronic myelogenous leukemia. The DNA-binding activity of the ubiquitously expressed ABL1 tyrosine kinase is regulated by CDC2-mediated phosphorylation, suggesting a cell cycle function for ABL1. The ABL1 gene is expressed as either a 6- or 7-kb mRNA transcript, with alternatively spliced first exons spliced to the common exons 2-11. [provided by RefSeq].	<b>Isotype:</b> IgG <b>SWISS:</b> P00519	<b>Applications:</b> Flow-Cyt (1ug/Test) <b>Reactivity:</b> Human (predicted: Mouse, Rat, Rabbit, Pig, Cow, Chicken, Dog, GuineaPig, Horse) <b>Predicted MW.:</b> 124 kDa <b>Subcellular Location:</b> Cell membrane ,Cytoplasm ,Nucleus
--	---	---

**— VALIDATION IMAGES —**

Blank control (black line) :Hela. Primary Antibody (green line): Rabbit Anti-c-Abl antibody ( bs-0583R ) Dilution:1ug/Test; Secondary Antibody (white blue line) : Goat anti-rabbit IgG-AF488 Dilution: 0.5ug/Test. Isotype control (orange line) : Normal Rabbit IgG Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.