bsm-41663M

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

ELISA (1:5000-10000)

Applications: WB (1:500-2000)

Reactivity: Human

Predicted 18 kDa

Subcellular Location: Secreted

MW.:

VEGFA/VEGF165 Mouse mAb

DATASHEET -

Host: Mouse Isotype: IgG2b Clonality: Monoclonal CloneNo.: 19D2 **GeneID: 7422 SWISS:** P15692

Target: VEGFA/VEGF165

Immunogen: Recombinant human VEGF165 Protein: 27-191/191.

Purification: affinity purified by Protein A

Storage: Size: 100ug

0.01M PBS (pH7.4). Size: 200ug (PBS only)

0.01M PBS

freeze/thaw cycles.


Background: Vascular endothelial growth factor (VEGF), originally known as

vascular permeability factor (VPF), is a signal protein produced by cells that stimulates the formation of blood vessels. To be specific, VEGF is a sub-family of growth factors, the platelet-derived growth factor family of cystine-knot growth factors. They are important signaling proteins involved in both vasculogenesis (the de novo formation of the embryonic circulatory system) and angiogenesis (the

growth of blood vessels from pre-existing vasculature).

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

- VALIDATION IMAGES -



25 ug total protein per lane of various lysates (see on figure) probed with VEGFA/VEGF165 monoclonal antibody, unconjugated (bsm-41663M) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.